

REFERENCES

HYPERTHYROIDISM CLINICAL CORRELATIONS

Bahn, R.S., Heufelder, A.E. (1993). Pathogenesis of Graves' ophthalmopathy. New Engl J Med, 329: 1468-1475.

Burch, H.B., Gorman, C.A., Bahn, R.S., Garrity, J.A. (1996). In Braverman, L.E., Utiger, R.D., (Eds.) Werner and Ingbar's The Thyroid – A Fundamental and Clinical Text, (7th ed.):536-553.

Burch, H.B., Wartofsky, L. (1993). Graves' ophthalmopathy: current concepts regarding pathogenesis and management. Endoc Rev, 14:747-793.

Cherner, R. (1997). Graves' disease: five questions physicians often ask. Consultant, 37 (10): 2688-2694.

Cooper, D. S. (1998). Antithyroid drugs for the treatment of hyperthyroidism caused by Graves' disease. Endocrinology and Metabolism Clinics of North America, 27 (1):225-247.

DeGroot, L. J. (1997). Radioiodine and the immune system. Thyroid, 7:259-264.

Gladstone, G. J. (1998). Ophthalmologic aspects of thyroid related orbitopathy. Endocrinology and Metabolism Clinics of North America, 27 (1): 91-100.

Haddad, G., (1998). Is it hyperthyroidism? You can't always tell from the clinical picture. Symposium: first of three articles on endocrine problems. Postgraduate Medicine, 104 (1):42-44, 53-55, 59.

Hashizume, K., Ichikawa, I., Sakurai, A. (1991). Administration of thyroxine in treated Graves' disease. N Engl J Med, 324: 947

<http://www.tfa.org/pub/tfa>

<http://ngdf.org/>

<http://mdconsult.com>

Kaplan, M.M., Meier, D. A., Dworkin, H. J. (1998). Treatment of hyperthyroidism with radioactive iodine. Endocrinology and Metabolism Clinics of North America, 27 (1), 205-223.

Madison, L.D. (1998). Today's approach to managing hyperthyroidism. Women's Health in Primary Care, 1 (5):452-465.

Marcocci, C., Bartelena, L., Bogazzi, F. (1992). Relationship between Graves' ophthalmopathy and type of treatment of Graves' hyperthyroidism. Thyroid, 2:171-178.

McIver, B., Rae, P., Beckett, G. (1996). Lack of effect of thyroxine in patients with Graves' hyperthyroidism who are treated with an antithyroid drug. N Engl J Med, 334:220.

McIver, B., Morris, J. (1998). The pathogenesis of Graves' disease. Endocrinology and Metabolism Clinics of North America, 27 (1):73-89.

Pfeilschifter, J., Ziegler, R. (1997). Suppression of serum thyrotropin with thyroxine in patients with Graves' disease: effects on recurrence of hyperthyroidism and thyroid volume. Eur J Endocrinol, 136: 81.

Reasner, C.A., Talbert, R. L. (2000). Thyroid disorders. In Dipiro, J.T., Talbert, R.L., Yee, G.C., Matzke, G. R., Wells, B.G., Posey, L. M. (Eds.) Pharmacotherapy: a pathophysiologic approach, (4th ed), Appleton and Lange. Stamford, Connecticut: 1244-1256.

Ringel, M.D., Cooper, D. S. (1996). Detection and management of hyperthyroidism: special considerations. Hospital Medicine, 32 (6): 15-20, 44-46.

Ryan, J. A., (1999). Severe neutropenia as an adverse effect of methimazole in the treatment of hyperthyroidism. Clinical Excellence for Nurse Practitioners, 3 (1): 2-6.

Schilling, J.S. (1997). Hyperthyroidism: diagnosis and management of Graves' disease. Nurse Practitioner, 22 (6): 72, 74-75, 83, 87-90.

Wells, B. G., Dipiro, J. T., Schwinghammer, T.L., Hamilton, C.W. (2000). Pharmacotherapy handbook, (2nd ed.). Stamford, Connecticut: Appleton and Lange, 216.